arXiv:hep-th/0310036v1 4 Oct 2003

Riemann-Silberstein representation of the complete Maxwell equations set

M. V. Cheremisin
A.F.Ioffe Physical-Technical Institute, St.Petersburg, Russia

(Dated: February 1, 2008)

Complete set of Maxwell equations is represented by single equation using the Riemann-Silberstein vector $\mathbf{F}=\mathbf{E}+i\mathbf{B}$. We demonstrate that the Fourier form of invariants E^2-B^2 and $\mathbf{EB}$ is proportional to dissipated power and equal to zero respectively.

PACS numbers: 03.50.-z, 03.50.De

The relative orientation of the magnetic and electric fields in a particular co-ordinate system is of great importance since both the real and the imaginary parts of the product $\mathbf{F}^2=E^2-B^2+2i\mathbf{EB}$ remain invariants along upon the Lorentz relativistic transformation. In this paper we represent Maxwell equations set by a single equation using Riemann-Silberstein [1], [2] formalism. Then, Fourier analysis allows us to eliminate the role playing by both invariants.

We use the set of conventional Maxwell equations

$$\mathrm{rot}\mathbf{B}=\frac{1}{c}\frac{\partial\mathbf{E}}{\partial t}+\frac{4\pi}{c}\mathbf{J}\qquad \mathrm{div}\mathbf{B}=0 \tag{1}$$
$$\mathrm{rot}\mathbf{E}=-\frac{1}{c}\frac{\partial\mathbf{B}}{\partial t}\qquad \mathrm{div}\mathbf{E}=4\pi\rho.$$

Here, the total current $\mathbf{J}=\mathbf{J}^{\mathrm{free}}+c\cdot\mathrm{rot}\mathbf{M}+\frac{\partial\mathbf{P}}{\partial t}$ and charge density $\rho=\rho^{\mathrm{free}}-\mathrm{div}\mathbf{P}$ are modified with respect to those for free carriers $\mathbf{J}^{\mathrm{free}},\rho^{\mathrm{free}}$ taking into account the polarization $\mathbf{P}$ and magnetization $\mathbf{M}$ of the system [3].

With the help of conventional definition $\mathbf{B}=\mathrm{rot}\mathbf{A}$, $\mathbf{E}=-\frac{1}{c}\frac{\partial\mathbf{A}}{\partial t}-\nabla\phi$ widely used to represent vector and electrical auxiliary potentials respectively Eqs.(1) yield

$$\Box\mathbf{A}=-\frac{4\pi}{c}\mathbf{J}+\nabla\psi\qquad \Box\phi=-4\pi\rho-\frac{1}{c}\frac{\partial\psi}{\partial t}, \tag{2}$$

where $\psi=\mathrm{div}\mathbf{A}+\frac{1}{c}\frac{\partial\phi}{\partial t}$, $\Box$ is the d'Lambert operator. We argue that Eqs.(2) provide the gauge transformation $\mathbf{A}\rightarrow\mathbf{A}+\nabla f,\phi\rightarrow\phi-\frac{1}{c}\frac{\partial f}{\partial t}$ is already satisfied.

It is to be noted that Eqs.(2) differ with respect to those known in literature. Usually, upon the extra condition $\psi=0$ one can readily recover the conventional Lorentz gauge [4] $\mathrm{div}\mathbf{A}+\frac{1}{c}\frac{\partial\phi}{\partial t}=0$, therefore Eqs.(2) become uncoupled. One can check, however, that the gauge transformation(see above), if applied, results in additional equation $\Box f=0$, which is always disregarded [5].

We now represent Maxwell equations by means of Riemann-Silberstein vector $\mathbf{F}$. Eqs.(1) yield

$$\mathrm{rot}\mathbf{F}=\frac{i}{c}(\dot{\mathbf{F}}+4\pi\mathbf{J}),\qquad \mathrm{div}\mathbf{F}=4\pi\rho \tag{3}$$

and finally

$$\Box\mathbf{F}=\dot{\mathbf{I}}-ic\cdot\mathrm{rot}\mathbf{I}, \tag{4}$$

where we introduce auxiliary vector $\dot{\mathbf{I}}=4\pi(\dot{\mathbf{J}}/c^2+\nabla\rho)$. One can complete the transformed Maxwell equations set implied above using the conventional boundary conditions as follows

$$F_{2n}-F_{1n}=4\pi\sigma,\qquad [\mathbf{nF_2}]-[\mathbf{nF_1}]=\frac{4\pi i}{c}\mathbf{j}, \tag{5}$$

where $F_{1,2}$ is the complex field above(below) the interface, $\mathbf{n}$ the normal to the interface. Then, we introduce the unit area related surface charge density $\sigma=\sigma^{free}-(P_{2n}-P_{1n})$ and surface current density $\mathbf{j}=\mathbf{j}^{free}+c([\mathbf{nM_2}]-[\mathbf{nM_1}])$.

The advantage of the Eqs.(4),(5) is, in fact, the compact representation for both the current(charge) sources of the electromagnetic field. In particular case of electric(magnetic) dipole irradiation Eq.(4) can reproduce, for example, the conventional result [6] as follows

$$\Box\mathbf{F}=-\frac{4\pi i}{c}\mathrm{rot}(\dot{\mathbf{K}}-ic\cdot\mathrm{rot}\mathbf{K})-4\pi\Box\mathbf{P}. \tag{6}$$

Here, we neglect free charge(current) densities, and, then introduce the complex electric(magnetic) dipole moment of the system $\mathbf{K}=\mathbf{P}+i\mathbf{M}$.

With the complex field $\mathbf{F}$ specified above the energy conservation law may be represented as follows

$$\frac{1}{8\pi}\frac{\partial}{\partial t}(\mathbf{FF}^*)=-\mathrm{div}\mathbf{S}-\mathbf{J}Re\mathbf{F}, \tag{7}$$

where $\mathbf{F}^*=\mathbf{E}-i\mathbf{B}$, $\mathbf{S}=\frac{ic}{8\pi}[\mathbf{FF}^*]=\frac{c}{4\pi}[\mathbf{EB}]$ is the conventional Poynting energy flux density.

Within Fourier formalism $\mathbf{F}=\int\limits_{-\infty}^{+\infty}\mathbf{F}_{k,\omega}e^{i\omega t-i\mathbf{kr}}\frac{d^3\mathbf{k}}{(2\pi)^3}\frac{d\omega}{2\pi}$ Eq.(4) yields

$$\mathbf{F}_{k,\omega}=\frac{c}{\kappa^2}[\mathbf{kI}_{k,\omega}]-\frac{i\omega}{\kappa^2}\mathbf{I}_{k,\omega}, \tag{8}$$

where $\mathbf{I}_{k,\omega}=4\pi(\frac{\mathbf{J}_{k,\omega}}{c^2}-\frac{\rho_{k,\omega}\mathbf{k}}{\omega})$, then $\kappa^2=k^2-\omega^2/c^2$. Since $(\mathbf{I}_{k,\omega}[\mathbf{kI}_{k,\omega}])=0$, one obtains

$$\mathrm{Re}\mathbf{F}^2_{k,\omega}=\frac{4\pi}{\kappa^2}(\mathbf{J}_{k,\omega}\mathbf{I}_{k,\omega}),\qquad \mathrm{Im}\mathbf{F}^2_{k,\omega}=0. \tag{9}$$

The latter result, namely that $(\mathbf{E}_{k,\omega}\mathbf{B}_{k,\omega})=0$ can be easily derived using Eqs.(1) as well. It's worthwhile to mention that $\mathbf{E}_{k,\omega} = -\frac{i\omega}{\kappa^2}\mathbf{I}_{k,\omega}$, and thus $\mathrm{Re}\mathbf{F}^2_{k,\omega}$ is proportional to dissipated energy $(\mathbf{J}_{k,\omega}\mathbf{E}_{k,\omega})$. For $\rho = 0$ Eqs.(9) correspond to electromagnetic wave propagating in vacuum when $\mathbf{EB} = 0, E^2 - B^2 = 0$.

In conclusion, we represent complete set of Maxwell equations by single equation using Riemann-Silberstein formalism. Fourier form of invariants $E^2 - B^2$ and $\mathbf{EB}$ is proportional to dissipated power and equal to zero respectively.

This work was supported by RFBR (grant 03-02-17588) and LSF(HPRI-CT-2001-00114)

[1] H.Weber, Die partiellen Differential-Gleichungen der Mathematischen Physik nach Riemann's Vorlesungen (Friedrich Vieweg und Sohn, Braunschweig), 348, 1901

[2] L.Silberstein, Ann. d Phys.(Leipzig) **22**, 579 1907; **24**, 783, 1907

[3] I.E.Tamm, Osnovi teorii elektrichestva, Moskow, 1957

[4] H.A.Lorentz, Encyklopedie der mathematischen Wissenschaffen, **5**, 156, 1903

[5] J.A.Stratton, Electromagnetic theory, New York, 1941

[6] L.D.Landau, E.M.Lifshits, The classical theory of fields, Pergamon, New York,1966